



NEONATAL MEDICATION GUIDELINE

PNEUMOCOCCAL CONJUGATE VACCINE, 13-VALENT (Prevenar®)

Scope (Staff): Nursing, Medical and Pharmacy Staff

Scope (Area): KEMH NICU, PCH NICU, NETS WA

This document should be read in conjunction with the [Disclaimer](#).

Quick Links

[Dose](#)

[Preparation & Administration](#)

[Side Effects & Interactions](#)

[Monitoring](#)

Restrictions

[Formulary: Unrestricted](#)

Description

Pneumococcal conjugate vaccine, 13-valent

Presentation

Suspension in pre-filled Syringe: 0.5mL

Storage

Refrigerate – do not freeze

Indication

Parent/Guardian consent is to be obtained prior to administration of all vaccinations.

Prevention of invasive diseases caused by *Streptococcus pneumoniae* in infants at:

- 2 months of age, and;
- 4 months of age, and;
- 12 months of age.

Additional dose **at 6 months** of age is recommended for:

- preterm infants born less than 28 weeks gestation;
- for Aboriginal and Torres Strait Islander infants;
- patients with Medical Risk Conditions – refer to [Australian Immunisation Handbook](#)

Dose

Intramuscular:

0.5mL

Administration

Intramuscular Injection ONLY

Shake syringe vigorously immediately prior to use to obtain a homogenous, white suspension.

Do not use if not uniformly suspended, or discoloured.

Inject 0.5 mL of suspension by intramuscular injection (IMI) to the anterolateral aspect of the thigh (slowly to reduce pain).

Pneumococcal Vaccine can be given at the same time as combined DTPa-hepB-IPV-Hib.

Pneumococcal Vaccine should be administered in the opposite leg to DTPa-hepB-IPV-Hib.

Compatible Fluids

Do not combine with any fluids or other vaccinations

Side Effects

Common: transient injection site reactions (pain, redness, itching, swelling or burning, small hard lump that may persist for some weeks), transient fever.

Serious: allergic reaction including anaphylaxis.

Monitoring

Infants receiving pneumococcal immunisation are to have a full set of observations taken prior to immunisations.

Continuous cardiac monitoring for 48 hours following immunisation.

If a transient fever occurs, Paracetamol may need to be administered.

Comments

Do not give during febrile illness or acute infection.

Parent/Guardian consent is to be obtained prior to administration of all vaccinations

Interactions

Co-administration of 13vPCV with Menactra (quadrivalent MenACWY vaccine) should be avoided. This is because Menactra may interfere with the immune response against some pneumococcal serotypes. Two other brands of MenACWY vaccine, Menveo or Nimenrix, are available and can be co-administered with

13vPCV without such interference. If 13vPCV and Menactra are inadvertently co-administered, a repeat dose of either vaccine is not needed.

Related Policies, Procedures & Guidelines

[WA Immunisation Schedule](#)

[Australian Immunisation Handbook – Pneumococcal disease](#)

[Immunisations](#)









References

Australian Medicines Handbook. Pneumococcal vaccines. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2022 [cited 2022 Nov 10]. Available from: <https://amhonline.amh.net.au/>

Australian Technical Advisory Group on Immunisation (ATAGI). Australian Immunisation Handbook, Australian Government Department of Health and Aged Care, Canberra, 2022, [cited 2022 Nov 10]. Available from: [Pneumococcal disease | The Australian Immunisation Handbook \(health.gov.au\)](#)

Neomed Formularies. Pneumococcal Vaccine In: The Royal Hospital for Women [Internet]. [South Eastern Sydney, New South Wales;2021 [cited 2022 Nov 11]. Available from: <https://www.seslhd.health.nsw.gov.au/royal-hospital-for-women/neomed-formularies>

PNEUMOCOCCAL CONJUGATE VACCINE, 13-VALENT (Prevenar®)

Keywords	Pneumococcal, prevenar, vaccine, immunisations, aboriginal				
Document Owner:	Head of Department - Neonatology				
Author/ Reviewer	KEMH & PCH Pharmacy/Neonatology Directorate				
Version Info:	4.0				
Date First Issued:	01/10/2001	Last Reviewed:	21/11/2022	Review Date:	21/11/2025
Endorsed by:	Neonatal Directorate Management Group			Date:	05/01/2023
NSQHS Standards Applicable:	<input checked="" type="checkbox"/>  Std 1: Clinical Governance <input type="checkbox"/>  Std 2: Partnering with Consumers <input type="checkbox"/>  Std 3: Preventing and Controlling Healthcare Associated Infection <input checked="" type="checkbox"/>  Std 4: Medication Safety		<input type="checkbox"/>  Std 5: Comprehensive Care <input type="checkbox"/>  Std 6: Communicating for Safety <input type="checkbox"/>  Std 7: Blood Management <input type="checkbox"/>  Std 8: Recognising and Responding to Acute Deterioration		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from WNHS HealthPoint.					

The health impact upon Aboriginal people has been considered, and where relevant incorporated and appropriately addressed in the development of this document (insert ISD Number). (Please refer to the Aboriginal Health Impact Statement and Declaration for Department of Health and Health Service Provider Guidelines – please delete once you have completed this).

This document can be made available in alternative formats on request for a person with a disability.

© Women and Newborn Health Service 2020

Copyright of this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.