



CLINICAL PRACTICE GUIDELINE

Infections: Antibiotic prophylaxis / medication: Hysterosalpingogram for infertility

This document should be read in conjunction with this [Disclaimer](#)

On the morning of the procedure commence:

- 50mg diclofenac every 8 hours (take with or soon after food). Maximum duration of 3-5 days.
- If dilated tubes or history of PID:
 - Doxycycline 100mg every 12 hours for 5 days
 - Metronidazole 400mg stat

References

1. Clifford V, Daley A. Antibiotic prophylaxis in obstetric and gynaecological procedures: A review. **ANZCOG**. 2012. 52(5) 412-9.

Related WNHS policies, procedures and guidelines

Obstetrics & Gynaecology: Infections: Antibiotic Prophylaxis and Treatment guidelines

Keywords:	hysterosalpingogram, infertility, AMS, antibiotic prophylaxis, gynae infection, antibiotics		
Document owner:	Obstetrics, Gynaecology & Imaging Directorate		
Author / Reviewer:	HoD Microbiology		
Date first issued:	June 2011	Version	3.0
Reviewed dates:	March 2015; Aug 2018	Next review date:	Aug 2021
Supersedes:	History: Originally located in Pharmacy (P4.14 then P3.9) then moved to O&G: Infections (A14) Supersedes: Version last reviewed March 2015		
Endorsed by:	GSMSC	Date:	02/08/2018
NSQHS Standards (v2) applicable:	3 Preventing and Controlling Infection, 4 Medication Safety		

Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.