



**CLINICAL PRACTICE GUIDELINE**

**Multiple pregnancy**

This document should be read in conjunction with the [Disclaimer](#)

**Contents**

**Antenatal management .....2**

    Background information ..... 2

    Counselling and management ..... 2

        Planned vaginal twin birth ..... 2

        Previous caesarean birth ..... 2

        Transverse lie of the presenting twin ..... 2

        Breech presentation of the first twin ..... 3

    Antenatal care ..... 3

        Routine antenatal investigation ..... 3

        Referrals ..... 3

        Frequency of antenatal clinic visits ..... 3

        Genetic screening ..... 4

        Fetal surveillance ..... 4

        Diet and nutrition ..... 5

        Hypertension ..... 5

        Parent education ..... 5

    Timing and mode of birth ..... 5

**Intrapartum management .....6**

    Admission ..... 6

    Intrapartum care ..... 6

        Intravenous access ..... 6

        Fetal monitoring ..... 6

        Analgesia ..... 7

        Diet and nutrition ..... 7

    Preparation for birth ..... 7

    Birth ..... 7

        Delivery of the first twin ..... 7

        Delivery of the second twin ..... 8

    Third stage management ..... 8

**References .....9**

# Antenatal management

## Background information

Multiple pregnancy is associated with increased risk of perinatal morbidity and mortality.<sup>1, 2</sup> Multiple gestation pregnancies are more likely to have complications of preterm labour, preterm premature rupture of the membranes, pre-eclampsia, pyelonephritis, and postpartum haemorrhage. Gestational diabetes is higher in twin and triplet pregnancies compared to singleton pregnancy.<sup>2</sup>

Early birth at 37 weeks gestation compared to ongoing expectant management for uncomplicated twin pregnancy is not associated with an increased risk of harm.<sup>3</sup>

Twin pregnancies are commonly divided according to zygosity or chorionicity as these have important implications for pregnancy and infant outcome.

## Counselling and management

Planned management and mode of delivery should be documented on the MR004 Obstetric Special Instruction Sheet after medical counselling with the mother.

### **Planned vaginal twin birth**

Birth of the healthy first twin presenting by vertex is associated with a low perinatal mortality and morbidity. However, some retrospective studies suggest there may be increased perinatal mortality due to intrapartum anoxia in the second twin at term. A recent RCT (2013) showed that in twin pregnancies after 32<sup>+0</sup> weeks gestation, with the first twin cephalic, the risks of fetal/neonatal death or serious neonatal morbidity were not significantly increased or decreased with a planned caesarean compared with planned vaginal birth.<sup>4</sup>

### **Previous caesarean birth**

Limited data is available concerning vaginal birth after caesarean in twin births, and currently there are no published randomised controlled trials. Recent studies have shown a trial of labour with twins does not appear to increase maternal morbidity, and perinatal morbidity is uncommon in gestation more than 34 weeks.<sup>5</sup> Other studies have found similar risk of uterine rupture to a singleton trial of labour<sup>6</sup>, with no more likelihood failing a trial of labour or experiencing major morbidity.<sup>7</sup>

The decision for trial of vaginal birth (VBAC) should be made in conjunction with the Consultant and the woman.

### **Transverse lie of the presenting twin**

An elective caesarean is the recommended mode of delivery.<sup>8</sup>

## Breech presentation of the first twin

The recommended mode of birth is caesarean section.<sup>8,9</sup> Studies have found significantly more depressed Apgar scores when the presentation of the first twin is breech, weighs more than 1500g, and has a vaginal birth.<sup>9</sup> See Clinical Guidelines [Abnormalities of Lie / Presentation](#).

## Antenatal care

### Routine antenatal investigation

Management as for a singleton pregnancy

### Referrals

Refer to the Maternal fetal Medicine team (MFM) for the following

- Monochorionic monoamniotic twins
- Monochorionic monoamniotic triplets
- Monochorionic diamniotic triplets
- Dichorionic diamniotic triplets
- Chorionic villus sampling(CVS) or amniocentesis
- Structural or chromosome anomaly
- Single fetal death on monochorionic twins
- Suspected twin – twin transfusion syndrome (TTTS)
- Severe early onset fetal growth restriction

Offer genetic counselling prior to screening for aneuploidy with nuchal translucency (NT) measurement

### Frequency of antenatal clinic visits

Women with twin pregnancies without complications are seen:

- 4 weekly until 28 weeks gestation
- 2 weekly until 34 weeks gestation
- weekly from 34 weeks gestation

In women with a high risk multiple pregnancy, the frequency of care is individualised. The consultant must be involved in the decision.

- Women with uncomplicated monochorionic, triamniotic and dichorionic triamniotic triplet pregnancies should have at least 11 antenatal appointments. Combine the appointments with the scans from approximately 11 weeks to 13 weeks 6 days gestation and then at estimated gestations of 16,18, 20, 22, 24, 26, 28, 30, 32 and 34 weeks.<sup>10</sup>
- Women with uncomplicated trichorionic, triamniotic triplet pregnancies should have appointments combined with scans from approximately 11 weeks 0 days

to 13 weeks 6 days and then at 20, 24, 28, 32 and 34 weeks. Offer an additional appointment without a scan at 16 weeks.<sup>10</sup>

## Genetic screening

- Offer nuchal translucency in the first trimester. Serum screening tests are not as sensitive in multiple gestations mainly due to limited available data. Serum levels markers are masked due to analytes from the normal and abnormal fetus both entering the maternal serum, and are in effect averaged together.<sup>11</sup> In monochorionic twin pregnancy maternal serum values can be used as each fetus has the same risk of aneuploidy. The role of these serum values is less certain in dichorionic twin pregnancies
- Where a first trimester screening for Down's syndrome cannot be offered to a woman with a twin pregnancy, consider second trimester screening and explain the potential problems of such screening including the increased risk of pregnancy loss associated with double invasive testing. **Second trimester serum screening for Down's syndrome must not be used in triplet pregnancies.**

## Fetal surveillance

### Ultrasound

- Perform an ultrasound at 12-13 weeks gestation. This ultrasound identifies whether the twins are dichorionic or monochorionic.
- Perform an anatomy scan at 19 weeks gestation. Congenital malformations are more common in multiple than singleton pregnancies. The incidence of malformations are higher in monozygotic twins than dizygotic twins.<sup>1</sup>
- Arrange serial growth and well-being scans according to chorionicity and clinical concern. The frequency of antenatal scans are generally
  - Monochorionic twins: screen for growth discordancy and TTTS every 2 – 3 weeks from 16 – 19 weeks gestation and then 2 weekly until delivery. Consider a fetal echo at 22 – 24 weeks.
  - Uncomplicated dichorionic diamniotic twins – screen for growth 4 weekly from 26 weeks.
  - If a woman with a twin or triplet pregnancy presents after 14 weeks 0 days, determine chorionicity at the earliest opportunity by ultrasound.<sup>10</sup>
  - If the woman presents late in pregnancy manage the pregnancy as monochorionic until proven otherwise.<sup>10</sup>

### Cardiotocograph (CTG) monitoring

- CTG monitoring should be done for
  - Discordant growth – weekly after 34 weeks. Serial ultrasounds provide the best method for monitoring discordant growth, with evaluation of fetal

wellbeing by use of a CTG monitoring, biophysical profiles and Doppler studies.<sup>12</sup>

- Risk factors of fetal compromise.

## Diet and nutrition

- Offer referral to Dietician Services. Multiple pregnancy increases calorie, protein, mineral and vitamin requirements.
- Recommend twice daily iron and folic acid supplementation. The risk for anaemia increases in multiple pregnancies.
- Recommend multivitamin supplementation for women with poor nutritional status. Iron deficiency anaemia is associated with pre-term delivery and low ferritin levels are linked to prematurity.

## Hypertension

- Advise women with twin and triplet pregnancies that they should take 75mg of aspirin daily from 12 weeks until the birth of the babies if they have one or more of the following risk factors for hypertension:
  - First pregnancy.
  - Age 40 years or older.
  - Pregnancy interval of more than 10 years.
  - BMI of 35 kg/m<sup>2</sup> at first visit.
  - Family history of pre-eclampsia.

## Parent education

- Advise the Parent Education department of all women with multiple pregnancies. This allows individual contact to provide information on specific classes and links to specialised community groups and services<sup>13</sup>.
- Consider booking Parent education classes early. Multiple gestation pregnancy has a higher risk for pre-term birth<sup>13</sup>.

## Timing and mode of birth

- The optimal timing of birth is uncertain with clinical support for both elective delivery at 37 weeks gestation (either by induction of labour or caesarean section) and waiting for labour to start spontaneously.
- NICE<sup>10</sup> recommends the following:
  - Monochorionic twin pregnancies – elective birth from 36<sup>+0</sup> weeks gestation after a course of prophylactic corticosteroids has been offered.
  - Dichorionic twin pregnancies – elective birth from 37<sup>+0</sup> weeks gestation.

- When appropriate obstetric experience is available, vaginal birth is the preferred mode of birth for all twin pregnancies that meet the following criteria<sup>4</sup>
  - Twins must be diamniotic
  - Twin 1 cephalic
  - Twin 2 is not >500g heavier than twin 1
  - Neither twin has any evidence of fetal compromise requiring caesarean section.
- Triplet pregnancies: elective birth from 35 weeks 0 days after a course of antenatal corticosteroids has been offered.<sup>10</sup>

## Intrapartum management

### Admission

If a woman presents in labour with a multiple pregnancy notify the:

- Midwifery Labour and Birth Suite Co-ordinator
- Consultant
- Obstetric Registrar
- Resident Medical Officer (RMO).

The Senior Registrar, Consultant and Anaesthetic Registrar should be advised of admission by the Obstetric Registrar.

### Intrapartum care

#### Intravenous access

- Insert an intravenous large bore cannula (16g). The risk for both intrapartum and postpartum haemorrhage is increased with multiple pregnancy.<sup>14</sup>
- Collect blood for:
  - Full blood picture
  - Group and hold
  - Cross-match blood if indicated e.g. anaemia

#### Fetal monitoring

Monitor the fetal heart rates (FHR) with the cardiotocograph (CTG) continuously in active labour (>4cm). Consider application of a fetal scalp electrode on twin one, and external monitoring on twin two if it is difficult to maintain continuous monitoring.<sup>14</sup>

## Analgesia

Discuss the option and benefits of intrapartum epidural analgesia. An epidural is recommended due to the increased risk of operative delivery in twin births, and the possibility of intrauterine manipulation of twin two.

## Diet and nutrition

Allow the woman to consume a low fat, low fibre, high calorie diet in labour. The incidence of aspiration pneumonia and Mendelson's Syndrome associated with emergency caesarean section in the developed world is relatively low<sup>15</sup>. Current evidence suggests that a policy of fasting in labour makes no difference to length of the labour or the obstetric or neonatal outcomes<sup>15</sup>.

## Preparation for birth

- Notify obstetric personnel to be present at the birth:
  - Registrar
  - Senior Registrar. If the senior obstetric registrar is not credentialed for twin birth, the Consultant should be present
  - Consultant if required.
- Notify the neonatal and paediatric staff to attend the birth.
- Ensure a portable ultrasound is available
- Advise the theatre coordinator and the duty Anaesthetic Registrar / Consultant that the birth is imminent.
- Ensure an oxytocin infusion is available to be used after the first twin is delivered. Oxytocin infusion may be required if uterine inertia occurs between twin births.<sup>14</sup>
- The oxytocin regime used between twins is:
  - 10 I.U. of oxytocin in 500mL of Hartmann's or Normal Saline commencing at 6mL/hour.

## Birth

### Delivery of the first twin

- Conduct the delivery of the first twin, if it is a cephalic presentation, as for a normal birth.
- Withhold the I.M. oxytocin after the birth of the first twin.
- Clamp and cut the umbilical cord after the birth of the first twin.
- Consider the commencement of an oxytocin infusion in consultation with the obstetric staff. This assists in the prevention of uterine inertia.<sup>14</sup> The solution

should be titrated according to the frequency/strength of contractions and consultation with medical staff.

### Delivery of the second twin

- Perform an abdominal palpation and vaginal examination immediately after delivery of twin one. This allows determination of the lie and presentation and position of twin two<sup>8</sup> and excludes cord presentation/prolapse.
- Confirm fetal presentation by portable ultrasound as required. External cephalic version or internal pedalic manipulation of the fetus may be required for malpresentation.
- Monitor the FHR of twin two continuously.
- Perform an artificial rupture of membranes (ARM) when clinically appropriate. The fetal presentation must be confirmed prior to performing an ARM. Cord presentation must be excluded.
- Withhold the third stage oxytocin until after delivery of the second twin.
- Aim to deliver the second twin within 30 minutes. If the condition of the second twin is satisfactory, the time factor becomes less important and should be weighed against the clinical situation.
- Collect cord blood from both twins after the birth of twin two.

### Third stage management

- Administer IM oxytocin after delivery of the second twin. See O&G clinical guideline: [Labour: Third Stage](#)
- Commence active management. A prophylactic oxytocin infusion is beneficial to prevent post-partum haemorrhage due to the risk of uterine atony which is increased with twin births<sup>9</sup>. See O&G clinical guideline: Postpartum Complications: Therapeutic and Prophylactic Oxytocin Infusion Regimens.



## References

1. Ayres A, Johnson TR. Management of Multiple Pregnancy: Prenatal Care - Part 1. **Obstetric and Gynecological Survey**. 2005;60(8):527-37.
2. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin Number 56. Multiple Gestation: Complicated Twin, Triplet, and High-Order Multifetal Pregnancy. **Obstetrics & Gynecology**. 2005;104(4):869-83.
3. Dodd JM, Deussen AR, Grivell RM, Crowther CA. Elective birth at 37 weeks' gestation for women with an uncomplicated twin pregnancy. **Cochrane Database of Systematic Reviews**. 2014 (2). Available from: <https://doi.org/10.1002/14651858.CD003582.pub2>
4. Barrett JFR, Hannah ME, Hutton EK, Willan AR, Allen AC, Armson BA, et al. A randomized trial of planned cesarean or vaginal delivery for twin pregnancy. **The New England Journal of Medicine**. 2013;369(14):1295-305. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24088091>
5. Varner MW, Leindecker S, Spong C, Et al. The Maternal-Fetal Medicine Unit cesarean registry: Trial of labor with a twin gestation. **American Journal of Obstetrics and Gynecology**. 2005;193:135-40.
6. Ford AA, Bateman BT, Simpson LL. Vaginal birth after casarean delivery in twin gestations: A large, nationwide sample of deliveries. **American Journal of Obstetrics and Gynecology**. 2006;195:1138-42.
7. Cahill A, Stamilio DM, Pare E, Et al. Vaginal birth after cesarean (VBAC) attempt in twin pregnancies: is it safe. **American Journal of Obstetrics and Gynecology**. 2005;193:1050-5.
8. Ayres A, Johnson TR. Management of Multiple Pregnancy: Labor and Delivery. **Obstetric and Gynecological Survey**. 2005;60(8):550-54.
9. Cruikshank DP. Intrapartum Management of Twin Gestations. **Obstetrics & Gynecology**. 2007;109(5):1167-76.
10. National Institute for Clinical Excellence (NICE). Multiple pregnancy: Antenatal care for twin and triplet pregnancies: CG 129: NICE; 2011 [cited 2019 April 1]. Available from: <https://www.nice.org.uk/guidance/cg129>
11. ACOG. ACOG Practice Bulletin Number 77 Screening for Fetal Chromosomal Abnormalities. **Obstetrics & Gynecology**. 2007;109(1):217-27.
12. Ayres A, Johnson TR. Management of multiple pregnancy: prenatal care--part II. **Obstet Gynecol Surv**. 2005;60(8):538-49.
13. Denton J, O'Brien W. Multiple pregnancy. In: Macdonald S, Johnson G, editors. *Mayes' midwifery*. Sydney: Elsevier; 2017. p. 965-78.
14. Dodd JM, Crowther CA. Evidence-based care of women with multiple pregnancy. **Best Practice & Research Clinical Obstetrics and Gynaecology**. 2005;19(1):131-53.
15. O'Sullivan G, Lui B, Shennan AH. Oral intake during labor. **International Anesthesiology Clinics**. 2007 45(1):133-47.

## Related WNHS policies, procedures and guidelines


KEMH Clinical Guidelines, Obstetrics & Gynaecology:

- [Abnormalities of Lie / Presentation](#)
- [Labour: Third Stage](#)
- [Postpartum Complications](#): Therapeutic and Prophylactic Oxytocin Infusion Regimens (available to WA Health employees through Healthpoint)

CMP Clinical Guideline: [Primary PPH at Home- Community Midwifery Program](#) (available to WA Health employees through Healthpoint)

## Useful resources (including related forms)

Forms: MR004 Obstetric Special Instruction Sheet

Keywords:	Multiple pregnancy, vaginal birth, caesarean section, breech presentation, antenatal clinics, fetal surveillance, monochorionic, monoamniotic, dichorionic, diamniotic, CTG, VBAC, ultrasound, triplet, twin, fetal, multiple pregnancy, fetal monitoring, intravenous access, analgesia, first twin, second twin, uterine atony, abruption, placenta, postpartum haemorrhage		
Document owner:	Obstetrics, Gynaecology & Imaging Directorate		
Author / Reviewer:	Evidence Based Clinical Guidelines Co-ordinator		
Date first issued:	June 2019 [New amalgamation]	Version	1.0
Reviewed:		Next review date:	June 2022
Supersedes:	<p><b>History:</b> In June 2019 amalgamated two individual guidelines on managing multiple pregnancies dating from March 2001.</p> <p><b>Supersedes:</b></p> <ol style="list-style-type: none"> <li>1. Multiple Pregnancy: Antenatal Management (dated Aug 2015)</li> <li>2. Multiple Pregnancy: Intrapartum Planned Vaginal Twin Birth (dated Aug 2015)</li> </ol>		
Endorsed by:	MSMSC (OOS approval- HoD Obstetrics, Coordinator of Midwifery, Midwifery Co-Director)	Date:	June 2019
NSQHS Standards (v2) applicable:	1  Governance		
<p><b>Printed or personally saved electronic copies of this document are considered uncontrolled.</b></p> <p><b>Access the current version from the WNHS website.</b></p>			