



CLINICAL PRACTICE GUIDELINE

Breech Presentation (Uncomplicated Term) – Planned Vaginal Birth

This document should be read in conjunction with the [Disclaimer](#)

Click on the heading and the hyperlink to take you to that section in the document.

[On admission – Management for women in labour](#)

[First stage management](#)

[Second stage management](#)

[Post birth management](#)

Background Information

Vaginal breech birth can be associated with a higher risk of perinatal mortality and short-term neonatal morbidity compared to birth by elective caesarean section¹, however study of long term follow-up at 2 years found that the neonatal neurological outcomes did not differ between either mode of birth even in the presence of serious short-term morbidity.²

Complications of vaginal breech birth include Erb's palsy, fractures to the clavicle, humerus or femur, and dislocation of the hips or shoulders. Trauma to the abdominal structures may occur if the fetal abdomen is grasped incorrectly, some bruising may be noted especially to male genitalia,³ and other complications such as cerebral haemorrhage or fractures, or spinal cord injury are additional risks.¹

Key Points

1. Planned term vaginal breech birth is a reasonable option provided there are no fetal or maternal contra-indications and the strict criteria is followed. The presence of an Obstetrician competent in breech delivery and facilities for immediate caesarean section are required⁴.
2. Women planning a breech birth who develop complications which are contraindications to a planned breech birth must be referred for review by the team consultant. If the consultant is unavailable or after hours, the woman must be reviewed in MFAU / Labour and Birth Suite by the Senior Registrar. The Consultant / Senior Registrar must have discussions with the woman and the junior medical staff.
3. The Consultant / Senior Registrar must have an informed discussion with the woman (and her support person if available) including options, recommendations and the possible outcomes.
4. This conversation and the final decision should be clearly documented in the notes by the medical officer with the appropriate level of seniority undertaking the counselling.

5. The Consultant Obstetrician is informed at the onset of labour, when the woman's cervix is fully dilated, and if there are concerns with maternal-fetal wellbeing or labour progress
6. The paediatric team is informed at the onset of labour, and should be present for the birth as per KEMH Clinical Guideline [Paediatric Team Attendance for 'At Risk' Births- LBS QRG](#)
7. Clinical pelvic examination should be performed to assess pelvic adequacy when assessing suitability for vaginal breech birth.
8. Induction of labour is not recommended and is considered non-standard management².
9. Augmentation of labour is not recommended but may be appropriate in the presence of uterine dystocia provided the consultant obstetrician is confident there is no fetopelvic disproportion^{1,2}.
10. Continuous cardiotocograph monitoring during labour should be performed⁴.
11. A vaginal examination should be performed if rupture of membranes occurs², or to confirm full dilatation prior to a woman pushing, ensuring she does not have a premature urge to push.
12. Cervical dilatation during *active* labour should occur at a rate of at least 1cm per hour.
13. The woman should not be encouraged to actively push until the breech has reached the pelvic floor and she has a strong urge, or the buttocks are on view.
14. If birth is not imminent after 1 hour of active pushing for a nullipara woman, or ½ hour for a multipara woman, a caesarean section should be initiated.
15. Breech extraction is not recommended during the breech birth of a singleton fetus.
16. Third stage oxytocic should not be administered until the fetal head is delivered.

Definition of an uncomplicated breech presentation

- Flexed or extended fetal legs.
- 37-42 weeks (women should be advised of risks associated with prolonged pregnancy).
- No evidence of cephalopelvic disproportion (CPD).
- Clinical estimation of the fetus >2.5kg or < 3.8kg.
- Well flexed head.
- No anticipated pelvic obstruction to birth.

On Admission – management for women in labour

1. Confirm the fetal presentation as flexed or extended breech of uncomplicated term breech and exclude contra-indications for vaginal breech birth by ultrasound.
2. Inform the Consultant Obstetrician.
3. Notify the paediatric team.
4. Commence cardiotocography (CTG) for continuous fetal heart rate monitoring.

5. Perform a digital vaginal examination to assess progress, and exclude cord presentation / prolapse.
6. Collect blood for a group and hold.

See the following two pages for planned vaginal breech management in the [first stage](#) and [second stage](#) of labour.

Flow Chart – Planned Term Singleton Vaginal Breech Birth

See KEMH Clinical Guideline [Breech Vaginal Birth – Quick Reference Guide](#)

First Stage Management

Care is the same as in cephalic presentation, with some additional care for the management of an uncomplicated term breech presentation.

Action	MANAGEMENT IN <u>FIRST STAGE</u>	ADDITIONAL INFORMATION
Monitoring labour progress	Cervical dilation should be 1cm per hour from 4cm for all women regardless of parity. In the absence of adequate progress in labour, caesarean section is recommended.	The Consultant Obstetrician should be advised of any delay in progress.
Augmentation	Not normally considered, however may be only used in individualised special circumstances for uterine dystocia ⁵ if there is no clinical suspicion of CPD. The decision for use is only made with consultant obstetrician approval.	Poor progress may be a risk factor for difficulty with delivery of the after coming head ⁶ . Intact membranes prevent risk for cord prolapse and artificial rupture of membranes is not recommended ³ .
Fetal Surveillance	Continuous CTG ⁴	
Vaginal Examination	<ul style="list-style-type: none"> • With spontaneous rupture of membranes • To confirm full dilatation if a woman has an urge to push. • Monitor routine progress of labour and more frequently as the situation requires. 	Excludes cord prolapse ⁷ . This confirms full dilatation of the cervix in the event of an urge to push.
Analgesia	Allow the women to choose. An epidural may be an option if the woman has a premature urge to push ⁸ .	
Bladder	Monitor 1-2 hourly	A full bladder may

Action	MANAGEMENT IN <u>FIRST STAGE</u>	ADDITIONAL INFORMATION
Management		impede descent of the breech.
Hydration	Fasting is not routinely required. Confirm medical recommendation.	Confirm with obstetric team hydration management.
Maternal Positioning	An upright position can be encouraged ⁸ .	An upright position may aid the descent of the breech ⁷ .
Additional Equipment	<ul style="list-style-type: none"> • Availability of the real time ultrasound machine 	
Medical notifications	Notify the Consultant Obstetrician: <ul style="list-style-type: none"> • At full dilatation • If poor progress of labour • If concerns of maternal-fetal wellbeing 	

Second Stage Management

Action	MANAGEMENT IN <u>SECOND STAGE</u>	ADDITIONAL INFORMATION
Confirm second stage	Perform a vaginal examination to confirm full dilatation prior to pushing.	Confirms that the woman is able to push if she has the urge.
Pushing	Encourage active pushing when the woman has a strong urge, or the buttocks are on view.	
Monitoring progress	Birth should be imminent after one hour of active pushing in a nullipara, and after ½ hour for a multipara. In the absence of adequate progress in second stage, caesarean section is recommended ⁶ .	The consultant obstetrician should be immediately notified of any delay in progress.
Fetal Surveillance	Continuous CTG ⁴	
Bladder Management	Consider urinary catheterisation prior to birth if the bladder is not emptied.	

Action	MANAGEMENT IN <u>SECOND STAGE</u>	ADDITIONAL INFORMATION
Position for birth	Dorsal or lithotomy	Following maternal consent the practitioner should utilise the maternal position with which they are familiar ⁶ .
Equipment	<ul style="list-style-type: none"> • Breech towel (warmed) • Lithotomy stirrups if necessary • Neville-Barnes' and Wrigley's Forceps immediately available 	
Analgesia	As indicated.	
Episiotomy	Not routine – should be performed when indicated to facilitate birth ⁶ .	
Birth principles	<ul style="list-style-type: none"> • No breech extraction • Traction/ fetal breech manoeuvres on breech are to be avoided unless necessary to expedite delivery of a partially expelled fetus in a timely fashion. • Gentle suprapubic pressure may aid flexion of the head⁶. • Do not handle / manipulate the cord. • Extended arms may be delivered by the Løvset manoeuvre.^{2, 8} Nuchal arms may be reduced with reverse Løvsets. • Aftercoming head may be delivered spontaneously, with forceps, or by the Mariceau-Smellie-Veit manoeuvre. • A small towel wrapped around the fetal hips is useful. 	<p>Can cause extension of the head and nuchal displacement of the arms.⁷</p> <p>May cause spasm of the cord⁸.</p> <p>Rapid birth of the head can cause sudden compression and risk for tentorium cerebelli tear.³</p> <p>Preserves warmth and provides a grip on the skin.</p>
Paediatrician	Contact the paediatric team to be present for the birth.	
Oxytocin for 3 rd stage	Withhold until the head is born.	

References and resources

1. Hannah ME, Hannah WJ, Hewson SA, et al. Planned caesarean section versus planned vaginal birth for breech presentation at term: A randomised multicentre trial. **The Lancet**. 2000;356:1375-83.
2. Society of Obstetricians and Gynaecologists of Canada. Vaginal delivery of breech presentation. **Journal of Obstetric Gynaecology of Canada**. 2009(June):557-66.
3. Thorogood C, Donaldson C. Disturbances in the rhythm of labour. **Midwifery preparation for practice**. 2nd ed. Sydney: Churchill Livingstones; 2010. p. 819-61.
4. Royal Australian and New Zealand College of Obstetricians and Gynaecologists (RANZCOG). **College Statement C Obs-11: Management of breech presentation at term** 2016. Available from: [https://www.ranzcog.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Management-of-breech-presentation-at-term-\(C-Obs-11\)-Review-July-2016.pdf?ext=.pdf](https://www.ranzcog.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Management-of-breech-presentation-at-term-(C-Obs-11)-Review-July-2016.pdf?ext=.pdf).
5. Society of Obstetrics and Gynecology of Canada. SGOC Clinical Practice Guideline No. 256. Substance Use in Pregnancy. **JOGC**. 2011(April):367-84.
6. Royal college of Obstetricians and Gynaecologists (RCOG). **Management of Breech Presentation (Green-top Guideline No. 20b)** 2017. Available from: <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg20b/>.
7. Lewis P. Malpositions and malpresentations. In: Macdonald S, Magill-Cuerden J, editors. **Mayer's Midwifery**. 14th ed. Sydney: Bailliere Tindall; 2011. p. 869-98.
8. Coates T. Malpositions of the occiput and malpresentations. In: Fraser DM, Cooper MA, editors. **Myles Textbook for Midwives**. 15th ed. Sydney: Churchill Livingstone; 2009. p. 573-605.

Related WNHS policies, procedures and guidelines

[Breech Presentation](#)

[Breech Presentation- Planned Vaginal Birth \(Quick Reference Guide\)](#)

Keywords:	breech vaginal birth, breech presentation, breech, uncomplicated term breech,		
Document owner:	Obstetrics, Gynaecology & Imaging Directorates		
Author / Reviewer:	O&G Evidence Based Clinical Guidelines		
Date first issued:	01/2012		
Last reviewed:	29/5/17	Next review date:	29/05/2020
Endorsed by:	Obs, Gynae & Imaging Directorate Management Committee	Date:	DDMonthYYYY

Standards Applicable:	NSQHS Standards: 1  Governance, 9  Clinical Deterioration,
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.	