

1 PRESCRIBING BLOOD COMPONENTS

Date Issued: December 2007
Date Revised: April 2011
Review Date: April 2013
Authorised by: KEMH Transfusion Committee
Review Team: KEMH Transfusion Committee

1.6 Refusal of Blood Products
Section 1
Transfusion Medicine Protocols
Women's & Newborn Health Services
Perth Western Australia

1.6 REFUSAL OF BLOOD PRODUCTS

Patient Choice includes the right to refuse blood components and blood products. This refusal may occur for a variety of reasons. In many instances this is a result of personal or religious beliefs. One of the groups who refuse blood products is the Jehovah's Witnesses. Jehovah's Witnesses' primary reason for declining blood transfusions is religious.

Refusal of consent for blood products must be clearly documented in the patient's medical file, including alternative means of treatment acceptable to the patient. However if a specific refusal of consent form for blood transfusion is available this should be used (KEMH Refusal form see below).

Jehovah's Witnesses WILL normally accept:


- Dextrose
- EPO (Recombinant Erythropoietin)
- G-CSF (granulocyte colony stimulating factor)
- Recombinant Factor VIIa (NovoSeven)
- Haemaccel
- Gelofusine
- Normal Saline, Hartmann's, Ringers
- Dextran

Jehovah's Witnesses WON'T normally accept:

- Whole Blood
- Red blood cells (packed cells)
- Platelets
- Fresh Frozen Plasma
- White blood cells (i.e. Granulocytes)
- Preoperative collection and storage of autologous blood.

Individual Jehovah's Witness decision may vary:

- Immunoglobulins (e.g. Anti D CMV etc.)
- Factor VIII (i.e. Biostate)
- Clotting factors (i.e. Prothrombinex)
- Albumin
- Cryoprecipitate
- Haemodilution
- Intraoperative Cell Salvage (if no allogeneic blood is used to prime the circuit and continuity is maintained with the vascular system.
- Post-Operative Reinfusion (i.e. Bellovac reinfusion and as above)



This guideline is for general INFORMATION PURPOSES ONLY. Each patient must be treated on an individual basis. One individual's wishes may not reflect all the above information.

For Management of obstetric patients the King Edwards Memorial Hospital Policy Guidelines shall be used, including the specific refusal of consent form.

http://wnhs.hdwa.health.wa.gov.au/_data/assets/pdf_file/0012/40017/Mr29599.pdf

KEMH Refusal Policy:

http://kemh.health.wa.gov.au/development/manuals/O&G_guidelines/sectionb/2/b2.14.pdf