

## 12 PLASMA DERIVED BLOOD COMPONENTS

Date Issued: January 2005  
Date Revised: July 2011  
Review Date: July 2013  
Authorised by: KEMH Transfusion Committee  
Review Team: KEMH Transfusion Committee

12.13 Intravenous Tetanus immunoglobulin  
Section 12  
Transfusion Medicine Protocols  
Women's & Newborn Health Services  
Perth Western Australia

### 12.13 INTRAVENOUS TETANUS IMMUNOGLOBULIN (4000IU)

#### DESCRIPTION

Each ampoule contains 4000 IU tetanus immunoglobulin for intravenous use.

#### INDICATIONS

Tetanus Immunoglobulin is available from the Blood Bank, to order please contact the Blood Bank Scientist and provide patient details and clinical history.

This product is for the management of established tetanus.

#### DOSE AND ADMINISTRATION

- One dose of 4000 IU administered by slow intravenous infusion will maintain minimum protective titre levels for at least 6 weeks

#### Further information may be found at

[http://www.csl.com.au/s1/cs/auhq/1217017237558/Web\\_Product\\_C/1196562715656/ProductDetail.htm](http://www.csl.com.au/s1/cs/auhq/1217017237558/Web_Product_C/1196562715656/ProductDetail.htm)