




CLINICAL PRACTICE GUIDELINE
NEWBORN EMERGENCY TRANSPORT SERVICE (NETS WA)

Resuscitation: Newborn

This document should be read in conjunction with the [Disclaimer](#)

Neonatal Resuscitation Algorithm – 2015 Update



WA Guidelines for newborn resuscitation

Adrenaline 1: 10 000 via ETT (1st dose; subsequent doses preferably by UVC)

- 1 mL for term infant
- 0.5 mL for preterm infant < 2kg or < 34 weeks gestation

If blender available commence oxygen for resuscitation at 30% and adjust as required.

Neopuff settings

- Term infant – 30 / 5 cmH₂O
- Preterm infant – 25 / 5 cmH₂O (< 2kg or < 34 weeks gestation)

Document owner:	Neonatal Directorate Management Committee		
Author / Reviewer:	Neonatal Directorate Management Committee		
Date first issued:	August 2009		
Last reviewed:	1 st October 2018	Next review date:	1 st October 2021
Endorsed by:	Neonatal Directorate Management Committee	Date endorsed:	23 rd October 2018
Standards Applicable:	NSQHS Standards: 1  Governance, 8  Acute Deterioration		
<p>Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.</p>			