



CLINICAL PRACTICE GUIDELINE

Labour: Partogram

This document should be read in conjunction with this [Disclaimer](#)

Contents

Key points	2
Documentation.....	2
Procedure	2
Admission & assessment findings plus medical/obstetric history.....	2
Maternal assessment.....	3
Fetal assessment.....	3
Labour assessment.....	4
Oxytocin administration.....	5
Investigations	5
Staff initials.....	5
Vaginal examination (VE)	5
Instrument/ pack count.....	7
Recording of staff.....	7
References.....	7

Key points

1. Alert and Action lines are drawn when the woman is in the active phase of labour.
2. The Alert line separates women into two groups, women with cervical dilatation:
 - equal to / greater than 1cm/hour.
 - slower than 1 cm/hour who are more likely to require an intervention.
3. The WHO partogram does not differentiate between nulliparous or multiparous women.⁴

Documentation

The partogram is a record of care, which constitutes a legal document but is also an avenue for identifying accountability in clinical practice. Therefore, accurate, legible and comprehensive entries should be made in accordance with guidelines and in black ink. Such entries should be made contemporaneously and authenticated with a full and legible signature.

Procedure

Admission & assessment findings plus medical/obstetric history

- Place the woman's identification label in the top left-hand corner.
- The admission details should be recorded as soon as practicable following arrival.
- Enter all details in the appropriate sections on the front of the partogram.

Date

- Record the commencement date at the top of the partogram. When the date changes at midnight write the new date above the times.

Time

- The numbered (0 1 2 3 etc) full vertical lines are hour lines.
- Note the exact time (e.g. 1450) of the first observation that you wish to record (e.g. fetal heart, vaginal examination, etc). Go back to the nearest whole hour – 1400 in this example.
- Fill in the time scale along the top of the partogram: 1400, 1500, 1600 etc (1400 is 0 hour line) and record the observation on the partogram (in this example just to the left of the 1500 line).
- Times must be written along the top of the partogram and may be written along the mid-section of the page.

Maternal assessment

- Record maternal blood pressure, pulse, temperature, respirations, and other observations (e.g. reflexes, blood sugar levels) on the graph at the top of the partogram.
- Using the measurements down the left side of the graph record:
 - Systolic blood pressure with a \wedge
 - Diastolic blood pressure with a \vee
 - Pulse with a \bullet

For example: 1400 1500 1600 1700 1800
 Observations at 1500 . . .
 and 1630hrs respectively

BLOOD	130						
			\wedge				
	120				\wedge		
\wedge							
\vee							
	110		\bullet				
	100					\bullet	
PULSE							
	90		\vee			\vee	
\bullet							
	80						
Temperature		37.1			37.9		
Respiration		22			20		
Reflexes							
Other (BSL/ SaO2)		3.1					

Fetal assessment

Record the FHR with a dot as follows:

- The vertical lines are half-hourly so that the quarter-hourly recordings can be made.
- During the second stage of labour the FHR is recorded 5 minutely in the boxes provided (if the fetus is not being continuously monitored). The time is written in the top left and the fetal heart rate is written below.
- Second stage fetal heart rate recordings must also be documented half hourly on the graph as in the first stage.

Amniotic fluid

- In the box that correlates with the correct time according to the vertical lines record half hourly absence or presence, and colour of fluid as follows:
 - I - membranes intact
 - C - amniotic fluid is clear
 - B - amniotic fluid is blood stained
 - M - amniotic fluid has meconium staining.

Labour assessment

Contractions

- These are recorded graphically below the fetal assessment information. An area of 5 blank vertical squares goes across the width of the graph and record contractions as 'frequency in 10 minutes'.
- Each square represents 1 contraction. Therefore if 2 contractions occur in 10 minutes, 2 squares will be shaded.
- Use the key for shading to demonstrate the strength of contraction.



Weak and/or 20-40 seconds duration



Moderate and 20-40 seconds duration



Strong and >40 seconds duration

Abdominal palpation

- Record in position abdominal palpation box.

Cervicograph

- The cervicograph is that section of the partogram which depicts cervical dilatation and descent of the presenting part in relation to time. Use of the cervicograph enables the progress of labour to be ascertained and delay in to progress readily recognized

Dilatation

- Record X for the cervical dilatation on the appropriate line and at the time the examination is carried out.

Descent

- Descent of the head is measured by abdominal palpation and is expressed in terms of fifths above the pelvic brim. Record O for the level of descent at each vaginal examination.

Alert line

- A line drawn from the point of cervical dilatation noted at the first vaginal examination in active labour. This line denotes a dilatation rate of 1cm/hour.

Action line

- A line parallel and 4 hours to the right of the alert line ¹.

Oxytocin administration

Units: record half hourly in black the number of units per 500mL of intravenous fluid.

mL/h: record in black the mL/h infusion rate.

For example:

		0900	1000	1100	1200		
Oxytocin	units		10	10	10	10	10
	mL/h		12 24	36	36	36	48

In this example the infusion commenced at 0930 at 12mL/h, increased to 24mL/h at 0945 and increased to 36mL/h at 1000.

Investigations

- Document any investigations performed since labour admission or those prior to induction of labour. For example: Full Blood Count, Group and Hold, vaginal swab.

Urine

- Record each void, measure the volume of urine as required.
- Record urinalysis (if required) for protein and ketones in the appropriate time box.

Staff initials

- Staff (including midwifery and medical students) that have performed observations are to provide their initials in the allocated boxes.
- Full name, signature, designation and initial are to be recorded on the reverse side of the partogram.

Vaginal examination (VE)

All VE's are to be documented on the partogram.

Date/time

- Record for each examination.

Indication

- Specify the reason for the vaginal examination.

Cervical effacement/length

- Estimate length in cm.

Dilatation

- Measured dilatation in cm.

Cervical position

- Stated as either anterior, posterior or midline.

Application

0: not applied

L: loosely applied

M: moderately applied

T: tightly applied.

Consistency

F: the cervix is firm to touch.

M: the cervix is medium to touch.

S: the cervix is soft to touch.

Membranes/liquor

- Use 'I', 'C', 'B', 'M'

Presentation

- Stated as cephalic, breech, shoulder, cord.

Position

- Record LOA, LOL, LOP, ROA, ROL, ROP.

Caput

0: no caput

+: small caput

++: moderate

+++: large caput

Moulding

0: no moulding

+: sutures are apposed

++: sutures overlapped but reducible

+++: sutures overlapped and not reducible.

Station

- This is measured in cms above (-) or below (+) the ischial spines. That is
 - Above spines: -5, -4, -3, -2, -1
 - 0 (at spines)
 - Below spines: +1, +2, +3, +4, +5

Head above brim (as per abdominal palpation)

- 5/5: completely above
- 4/5: sinciput high, occiput easily felt
- 3/5: sinciput easily felt, occiput felt
- 2/5: sinciput felt, occiput just felt
- 1/5: sinciput felt, occiput not felt
- 0/5: none of head palpable

FHR post-VE/FBS

- Record the fetal heart rate after a vaginal examination and the result of a fetal blood sample here.

Bishops score

- See scoring guide on the partogram. Record if induction of labour is required.

Performed by

- Include signature, printed name and designation.

Instrument/ pack count

- Record the number of birth instruments and/or packs utilized on the Instrument/Pack Count.
- If perineal repair is required document the Instrument and pack count on the record of THE repair.

Recording of staff

- Legibly document name, signature, designation and initials.

References

1. World Health Organisation. WHO recommendation on the use of active phase partograph with a four hour action line for monitoring the progress of labour. 2014.

Related legislation and policies



Related WNHS policies, procedures and guidelines

KEMH Clinical Guidelines:

- Labour and Birth: First stage; Second Stage

Useful resources (including related forms)

Partogram

Keywords:	partogram, cervical dilation, action line, alert line, labour		
Document owner:	Obstetrics, Gynaecology & Imaging Directorate		
Author / Reviewer:	CMC Labour & Birth Suite		
Date first issued:	Nov 2001		
Reviewed:	; Feb 2015; Oct 2018; (March 2019- minor amendment to table on page 5)	Next review date:	Oct 2021
Supersedes:	Version last reviewed Oct 2018		
Endorsed by:	MSMSC [with OOS A/HoD Obstetrics, CMC LBS; A/Coordinator of Midwifery]	Date:	27/11/2018
NSQHS Standards (v2) applicable:	1  Governance, 8  Recognising & Responding to Acute Deterioration		
<p>Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.</p>			