



NEONATAL MEDICATION GUIDELINE

Dexamethasone

Scope (Staff): Nursing, Medical and Pharmacy Staff

Scope (Area): KEMH NICU, PCH NICU, NETS WA

This document should be read in conjunction with the [Disclaimer](#).

Quick Links

[Dose](#)

[Preparation & Administration](#)

[Side Effects & Interactions](#)

[Monitoring](#)

Restrictions

[Formulary: Restricted](#)

Requires Neonatologist or relevant specialist review within 24 hours of initiation.

Description

Long acting synthetic glucocorticoid (adrenal steroid hormone), anti-inflammatory and immunosuppressive.

Presentation

Vial: 4mg/mL (4000microg/mL)

Oral solution: 1mg/mL (1000microg/mL)

Storage

Vial: Store at room temperature, below 25°C.

Oral solution: Refrigerate, do not freeze.

Dose

Facilitating extubation in ventilated babies with evolving or established chronic lung disease (Low dose DART Regimen)

IV/Oral:

Note - The weaning regime can be shortened or lengthened depending on the infant's clinical response.

Day	Dose	Frequency	Duration
Day 1-3	75 microgram/ kg/ dose	12 hourly	72 hours
Day 4-6	50 microgram/ kg/ dose	12 hourly	72 hours
Day 7-8	25 microgram/ kg/ dose	12 hourly	48 hours
Day 9-10	10 microgram/ kg/ dose	12 hourly	48 hours

Treatment of confirmed COVID-19 where the neonate requires oxygen and/or ventilation

IV/Oral:

Per Low dose DART regimen in table above.

Prevention of post intubation stridor in neonates who have had repeated, traumatic or prolonged intubation

IV/Oral:

250 microgram/kg/dose every 8 hours for a total of 3 doses.

Recommended to begin at least 4 to 12 hours prior to extubation.

For respiratory insufficiency and oedema with acute non-infectious laryngospasm

IV/Oral:

250 microgram/kg/dose every 8 hours for a total of 3 doses.

Dose Adjustment

Adjust dose according to response as per consultant advice.

Renal impairment: no dose adjustment documented.

Hepatic impairment: no dose adjustment documented.

Preparation

IV (diluted)

Available from CIVAS (*KEMH only*)

Withdraw 0.5mL (2mg) of dexamethasone and add 19.5mL of compatible fluid.

Concentration = 2mg/20mL = 2000microgram/20mL.

Final Concentration = 100microgram/mL.

Oral

Dilution required for doses less than 100micrograms:

Take 1mL (1000 micrograms) of oral dexamethasone solution and dilute to a final volume of 10mL with Water for Irrigation.

Concentration = 1000microgam/10mL = 100microgam/mL

Administration**IV Push (diluted)**

Slow push over 3-5 minutes

IV Infusion (undiluted)

For 250micrg/kg doses for post-intubation stridor only.

Use undiluted and infuse over 15 minutes.

Oral

For doses less than 100micrograms

Give with or immediately after feeds to minimise gastric irritation

Compatible Fluids

Sodium Chloride 0.9%, Glucose 5%

Y-Site Compatibility

Refer to KEMH Neonatal Medication Guideline: [Y-Site IV Compatibility in Neonates](#)

Side Effects

Common: hyperglycaemia, hypertension, behavioural disturbances, increase in urinary calcium excretion, sodium and water retention.

Serious: sepsis, masking of signs of infection, adrenal suppression, acute adrenal insufficiency in abrupt withdrawal, G.I. bleeding, osteoporosis, fractures, growth restriction, increased risk of cerebral palsy, delayed wound healing, skin atrophy, cushingoid appearance.

Interactions

Avoid concurrent use with Indomethacin for PDA treatment

Monitoring

Monitor blood glucose levels, blood pressure, electrolytes.

Bone bloods for long term therapy. Signs and symptoms of sepsis.

Related Policies, Procedures & Guidelines

CAHS COVID-19 Resources:

[Clinical Care of Paediatric Patients During the COVID-19 Pandemic](#)

Australian guidelines for the clinical care of people with COVID-19:

[Section 6.1.3.3: Corticosteroids for children and adolescents](#)

References

Ainsworth SB. Neonatal formulary 7 : drug use in pregnancy and the first year of life. Seventh ed. Chichester (West Sussex): John Wiley & Sons Inc.; 2015. p 175.

Takemoto CK, Hodding JH, Kraus DM. Pediatric & neonatal dosage handbook with international trade names index: a universal resource for clinicians treating pediatric and neonatal patients. 27th ed. Hudson (Ohio): Lexicomp; 2020-2021. 693.

Australian Medicines Handbook. Dexamethasone. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2020 [cited 2021 Jan 22]. Available from: <https://amhonline.amh.net.au/>

Truven Health Analytics. Title e.g. Dexamethasone. In: NeoFax [Internet]. Greenwood Village (CO): Truven Health Analytics; 2021[cited 2021 Jan 22]. Available from: <https://neofax.micromedexsolutions.com/>









Society of Hospital Pharmacists of Australia. Dexamethasone. In: Australian Injectable Drugs Handbook [Internet]. [St Leonards, New South Wales]: Health Communication Network; 2021 [cited 2021 Jan 22]. Available from: <http://aidh.hcn.com.au>

Plover C, Porrello E. Paediatric injectable guidelines 2019 ed. Flemington (Victoria): The Royal Children's Hospital Melbourne; 2019. p. 33

DART Regimen Paediatric Research. 2004 Sep; 56(3):477.

Australian National COVID-19 Clinical Evidence Taskforce. Corticosteroids for children and adolescents. In: Australian guidelines for the clinical care of people with COVID-19 [Internet]. Victoria; March 2022 [cited 2022 March 11]. Available from:

<https://app.magicapp.org/#/guideline/L4Q5An/section/EaD9Dn>

Keywords	Dexamethasone, DART, stridor, chronic lung disease, steroid, COVID-19, COVID				
Document Owner:	Head of Department - Neonatology				
Author/ Reviewer	KEMH & PCH Pharmacy/Neonatology Directorate				
Version Info:	July 2020 – full review (V3) January 2021 – full review (V3.1) March 2022 – added COVID indication and resources, updated template (V3.2)				
Date First Issued:	March 2001	Last Reviewed:	March 2022	Review Date:	January 2024
Endorsed by:	Neonatal Directorate Management Group			Date:	22/03/2022
NSQHS Standards Applicable:	<input checked="" type="checkbox"/>  Std 1: Clinical Governance <input type="checkbox"/>  Std 2: Partnering with Consumers <input type="checkbox"/>  Std 3: Preventing and Controlling Healthcare Associated Infection <input checked="" type="checkbox"/>  Std 4: Medication Safety		<input type="checkbox"/>  Std 5: Comprehensive Care <input type="checkbox"/>  Std 6: Communicating for Safety <input type="checkbox"/>  Std 7: Blood Management <input type="checkbox"/>  Std 8: Recognising and Responding to Acute Deterioration		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from WNHS HealthPoint.					

The health impact upon Aboriginal people has been considered, and where relevant incorporated and appropriately addressed in the development of this document (insert ISD Number). (Please refer to the Aboriginal Health Impact Statement and Declaration for Department of Health and Health Service Provider Guidelines – please delete once you have completed this).

This document can be made available in alternative formats on request for a person with a disability.

© Women and Newborn Health Service 2020

Copyright of this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.