



**OBSTETRICS AND GYNAECOLOGY
 CLINICAL PRACTICE GUIDELINE**

Oxygen therapy

Scope (Staff):	WNHS Obstetrics and Gynaecology Directorate staff
Scope (Area):	This guideline applies to specific Obstetrics and Gynaecology clinical areas of KEMH- Wards 3, 4, 5 and 6 only. Notes: For general information: Refer to Oxygen Therapy (SCGH/OPH) (available to WA Health employees through HealthPoint). For OPH WNS- see pregnancy section below and above SCGOPHCG guideline.
This document should be read in conjunction with this Disclaimer	

Contents

Key points2

Patients with severe chronic lung disease or other conditions at risk of hypercapnoeic failure3

Reduction and discontinuation of oxygen therapy3

Pregnancy3

Posture and oxygenation4

Oxygen delivery systems4

Documentation.....5

References6



Key points

1. No patient shall be denied oxygen therapy in an emergency or clinical deterioration event.¹ Patients in cardiac arrest and / or respiratory arrest shall be managed as per Clinical Guideline Obstetrics and Gynaecology: [Acute Deterioration \(Adult\): Resuscitation and Life Support](#).
2. All other patients with suspected or known tissue hypoxia shall have oxygen therapy initiated immediately by the attending health care professional (Doctor, Nurse, Midwife or Physiotherapist)
3. Patients commenced on acute oxygen therapy shall be examined by a doctor as soon as possible and if possible, initial investigations may include an arterial blood gas (ABG), haemoglobin level and chest X-ray.
4. **Observations:**
 - Initial and regular observations should include vital signs (respiratory rate, oxygen saturation, oxygen flow rate, heart rate, blood pressure, temperature and consciousness)
 - Patients receiving oxygen therapy shall be monitored by pulse oximetry at least as frequently as other observations and clearly documented on the patient's observation and response chart (ORC)¹ with the oxygen flow rate.
5. **Escalation and review of patients** must be managed as per the ORC for patients:
 - commenced on oxygen for deteriorating SpO₂¹
 - requiring increasing oxygen flow rates or decreasing saturation readings¹
 - developing other signs of deterioration (e.g. acute breathlessness, decreasing conscious state)
6. **If the patient cannot be stabilised:** Specialist medical advice shall be obtained promptly from the Anaesthetic Department.
7. Once the patient is stable, the oxygen therapy must be prescribed on the dedicated 'Oxygen Therapy Medication Chart' (MR 810.70) by a Medical Officer.
 - The MR 810.70 shall define the target oxygen saturation, oxygen therapy delivery device, range for oxygen flow or percent of inspired oxygen, and when oxygen is to be applied. The prescription shall be signed, dated and name of prescriber printed legibly. The prescription shall be reviewed **daily**.
8. On the Medication Chart (MR 810), document (tick) in "Additional Charts" – 'Other' and indicate there is an Oxygen Therapy Medication Chart in use.
9. Oxygen therapy shall be titrated to the lowest concentration that meets oxygenation goals:
 - 94-98% for all patients except those with, or at risk of, hypercapnoeic respiratory failure
 - 88-92% for those patients with or at risk of hypercapnoeic failure²

Patients with severe chronic lung disease or other conditions at risk of hypercapnoeic failure

Oxygen therapy shall be prescribed cautiously to patients with severe chronic lung disease and other conditions at risk of hypercapnoeic respiratory failure e.g. morbid obesity, neuromuscular disorders, and chest wall disorders.¹

1. These patients who require oxygen therapy shall be commenced on nasal prongs at 0.5 – 2L/min or oxygen using a Venturi mask at 24-28%.
2. Titrate oxygen to maintain an oxygen saturation level of 88-92% ¹ as per documented modification.
3. Regular monitoring of SpO₂ to maintain in the prescribed target range.¹
4. In patients with hypercapnoea, ABGs should be repeated one-hour post commencement of oxygen therapy and as per medical instruction.¹ Assisted ventilation (either non- invasive or intubation) should be considered in patients where respiratory acidaemia develops (pH < 7.35 with an increase in PaCO₂).
5. If nebulised bronchodilators are required, the nebuliser is best driven by compressed air with oxygen therapy given concurrently by nasal cannulae at 2 – 4L / min to maintain an oxygen saturation of 88-92%. If compressed air is not available, the nebuliser can be driven by oxygen at a flow rate of 6-10L / min for 3 - 4 minutes.

Reduction and discontinuation of oxygen therapy

- Oxygen therapy shall be reduced and discontinued in stable patients with satisfactory oxygen saturation. The oxygen saturation shall be above the target range or have been at the upper end of the target range for at least 4 hours.
- Discuss with the Medical Officer prior to reduction or ceasing of oxygen therapy in patients with severe chronic lung disease or other conditions at risk of hypercapnoeic failure.
- Oxygen therapy shall cease when the patient is able to maintain oxygen saturation in the target range when breathing room air. Oxygen saturation on room air shall be monitored for at least 5 minutes after discontinuing oxygen therapy and shall be rechecked at 1 hour. Oxygen therapy shall be recommenced if the oxygen saturation falls below the target range.
- **Once discontinued, the Nurse/Midwife is to document 'Ceased' on MR810.70.**

Pregnancy

- All women with hypoxaemia who are more than 20 weeks gestation should be managed with lateral tilt to improve cardiac output.
- The use of oxygen therapy during labour in normoxic women has been associated with acidosis of cord blood.

Posture and oxygenation

To increase oxygenation patients should be nursed in the upright position unless:

- the upright position causes discomfort
- immobilisation is required for a medical/surgical condition
- the patient is hypotensive
- the patient is recovering from a seizure

Oxygen delivery systems

For additional details on use oxygen delivery systems, safety, comparison of delivered oxygen, and reference ranges for ABGs, see SCGOPHCG Clinical Practice Guideline (CPG) [Oxygen Therapy](#) (available to WA Health employees through HealthPoint).

- Please note that the SCGH/OPH guideline is for clinical information only. Information contained in it regarding contacts and paperwork (forms) are not applicable for KEMH.
- KEMH documents on ORCs and Oxygen Therapy Medication Chart (MR810.70)
- OPH WNS uses [SCGOPHCG Oxygen Therapy](#) (available to WA Health employees through HealthPoint) and Oxygen Therapy Prescription Chart.

Simple nasal cannula

- Flow rate greater than 4 litres/minute oxygen, should not be used¹
- Refer to section “Nasal Cannula” within SCGOPHCG CPG [Oxygen Therapy](#) (available to WA Health employees through HealthPoint).

Humidified high flow nasal prong oxygen

- See sections “Humidification” and “Humidified Nasal Cannula” within [SCGOPHCG Oxygen Therapy](#) guideline (available to WA Health employees through HealthPoint).

Simple (Hudson) mask

- The flow should be 5L-10L/min because lower flows can cause rebreathing of exhaled CO₂.¹
- The mask is suitable for patients with hypoxaemic (Type 1) respiratory failure but not for patients with hypercapnoeic (Type II) respiratory failure
- Refer to section “Simple Face Mask” within [SCGOPHCG Oxygen Therapy](#).

Reservoir (non-rebreathing) mask

Reservoir masks can be used to provide a higher FiO_2 than simple masks. They are most suitable in an emergency (e.g. shock, trauma) where CO_2 retention is less relevant. The bag must be fully inflated before application to the patient and should remain inflated. Deflation suggests a leak or inadequate oxygen flow and may be a sign of deterioration.

- See section “Partial and Full Non-Rebreather Mask” within [SCGOPHCG Oxygen Therapy](#) guideline.

Graduated (e.g. Venturi) mask

Venturi masks provide a more predictable oxygen concentration to the patient. The masks are available in the following concentrations: 24%, 28%, 31%, 35%, 40% and 50% depending on the colour-coded mask attachment used. They are suitable for all patients needing a known concentration of oxygen. The 24% and 28% Venturi masks are particularly suited to those at risk of CO_2 retention. The oxygen flow required to achieve the appropriate concentration is defined on the colour coded mask attachment. The accuracy of the oxygen concentration delivered is greatly reduced if the mask is not accurately placed on the patient’s face, the flow is too low or the diluter jet is obstructed. The latter may be caused by clothing and can be prevented by the use of a protective hood over the jet diluter.

- See section “Graduated “Venturi” Type Mask” within [SCGOPHCG Oxygen Therapy](#).

Documentation

‘Oxygen Therapy Medication Chart’ (MR 810.70)

1. The Medical Officer orders the oxygen therapy on the oxygen prescription form. This is then attached to the medication chart
2. Document in the “Additional charts” sections of the Medication Chart to indicate the patient has been prescribed oxygen therapy
3. The Registered Nurse/Midwife signs the chart once per shift to indicate the oxygen prescription has been checked
4. The MR 810.70 is reviewed every 24 hours, or according to clinical state by the medical team
5. Changed oxygen therapy is ordered and documented on the MR 810.70
6. Record oxygen saturations on the relevant ORC
7. Record commencement, completion and changes to oxygen therapy on the ORC
8. At the beginning of the shift, the Midwife / Registered Nurse will check the order for oxygen therapy with the oxygen therapy the patient is receiving, and then document the type and flow rate of the oxygen on the ORC.

References

1. Sir Charles Gairdner Osborne Park Health Care Group [SCGOPHCG]. Clinical practice guideline: Oxygen therapy: SCGOPHCG. 2023. Available from: <https://healthpoint.hdwa.health.wa.gov.au/policies/Policies/NMAHS/SCGH/SCGOPHCG.CPG.OxygenTherapy.pdf>
2. Australian and New Zealand Committee on Resuscitation. ANZCOR Guideline 11.6.1: Targeted oxygen therapy in adult advanced life support: ANZCOR. 2016. Available from: <https://resus.org.au/guidelines/>

Related legislation and policies

SCGOPHCG CPG: [Oxygen Therapy](#) (WA Health employees access via HealthPoint)

Related WNHS policies, procedures and guidelines

WNHS Clinical Guideline, [Obstetrics and Gynaecology](#): Acute Deterioration (Adult): Resuscitation and Life Support

WNHS Policy: [Recognising and Responding to Acute Clinical Deterioration \(Physiological and Mental Health\)](#) (including code blue calling and escalation)

Useful resources (including related forms)

Forms

- Observation and Response Charts (ORC)
- Oxygen Therapy Medication Chart (MR810.70)

Keywords:	Oxygen therapy, oxygen administration		
Document owner:	Obstetrics and Gynaecology Directorate		
Author / Reviewer:	Consultant Staff Specialist Anaesthesia and Pain Management; Evidence Based Clinical Guidelines Co-ordinator OGD		
Date first issued:	June 2014(v1)	Version:	4.1
Reviewed dates:	(Jan 2015 amendment v1.2); Aug 2016(v2); Apr 2017(v2.1); Apr 2020(v3); Apr 2023(v4); 19/4/23 (v4.1-minor amendments)	Next review date:	Apr 2026
Endorsed by:	Obstetrics and Gynaecology Directorate Management Committee	Date:	05/04/2023
NSQHS Standards (v2) applicable:	<input checked="" type="checkbox"/>  1: Clinical Governance <input type="checkbox"/>  2: Partnering with Consumers <input type="checkbox"/>  3: Preventing and Controlling Healthcare Associated Infection <input checked="" type="checkbox"/>  4: Medication Safety	<input checked="" type="checkbox"/>  5: Comprehensive Care <input checked="" type="checkbox"/>  6: Communicating for Safety <input type="checkbox"/>  7: Blood Management <input checked="" type="checkbox"/>  8: Recognising and Responding to Acute Deterioration	
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from WNHS HealthPoint.			

Version history

Version number	Date	Summary
1	June 2014	First version. Archived- contact OGD Guideline Coordinator for previous versions. Original titled as A.12.1: 'Respiratory Support: Oxygen Therapy'
1.2	Jan 2015	Minor amendment- Removed numbers from title for alphabetical listing
2	Aug 2016	Lowest oxygenation goal for all patients except those with / at risk of hypercapnoeic respiratory failure raised to 95%
2.1	Apr 2017	<ul style="list-style-type: none"> • Content moved within guideline for clarity • Once discontinued, the Nurse/Midwife is to document 'Ceased' on the MR810.70
3.0	Apr 2020	<ul style="list-style-type: none"> • Monitor and escalate as per relevant ORC chart. • Lowest oxygenation goal lowered to 94% for all patients except those with, or at risk of, hypercapnoeic respiratory failure. ANZCOR referenced for target oxygen values. • Tick on Medication Chart if an Oxygen Chart is in use. Oxygen sticker no longer used in medication chart. • Patients at risk of hypercapnoeic failure- nasal prongs added as an initial option • Oxygen delivery systems- made more concise. Links to SCGH guideline for added details on how to use equipment
4	Apr 2023	Evidence on this topic was reviewed and overall guidance remains unchanged. Minor changes, links and formatting have been made. In hypercapnoeic patients, ABGs 1 hour post commencement
4.1	19/04/2023	Minor amendments- new SCGOPHCG clinical practice guideline (in place of nursing practice guideline)- hyperlinks updated. OPH no longer has separate oxygen guideline- uses SCGOPHCG guideline.

This document can be made available in alternative formats on request for a person with a disability.

© North Metropolitan Health Service 2023

Copyright to this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the *Copyright Act 1968*, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.

www.nmhs.health.wa.gov.au