ADULT Medication Monograph

BENZYLPENICILLIN

This document should be read in conjunction with this **DISCLAIMER**

Antimicrobial Restriction – Unrestricted

Presentation	Vial (Powder for reconstitution): 600mg, 3g					
Dose	Prevention of neonatal Group B streptococcal disease					
	IV:					
	Loading dose: 3g					
	Maintenance: 1.8g every 4 hours until birth					
	Tertiary Syphilis					
	IV:					
	1.8g every 4 hours for 15 days					
	Pregnancy and Congenital Syphilis					
	IV:					
	50mg/kg every 12 hours for 10 days					
Administration	IV injection:					
	Step 1 Reconstitution:					
	600mg vial: Reconstitute vial with 3.6mL WFI. Final concentration is 150mg/mL.					
	3g vial: Reconstitute vial with 13mL WFI. Final concentration is 200mg/mL.					
	Step 2 Administration:					
	Inject SLOWLY. Do NOT inject faster than 300mg/minute (e.g. 3g over 10 minutes) as rapid IV injection of large doses may cause seizures.					
	IV infusion:					
	Step 1 Reconstitution:					
	Reconstitute as above					
	Step 2 Dilution:					
	Dilute dose in 100mL of glucose 5% or sodium chloride 0.9%					

	Step 3 Administration:				
	Infuse over 30 minutes to 1 hour. Use immediately after preparation				
	<u> </u>				
Pregnancy	1 st Trimester: Safe to use				
	2 nd Trimester: Safe to use				
	3 rd Trimester: Safe to use				
Breastfeeding	Safe to use				
Clinical Guidelines	WNHS Policy: Antimicrobial Stewardship				
and Policies	KEMH Pharmaceutical & Medicines Management Guideline:				
	Medication Administration				
	KEMH Clinical Guideline: O&G: <u>Pre-labour Rupture of Membranes at Term</u>				
	KEMH Clinical Guideline: O&G: Group B Streptococcal Disease				
	KEMH Clinical Guideline: O&G: Sexually Transmitted Infections: Syphilis				
References	Australian Medicines Handbook. Benzylpenicillin. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2019 [cited 2019 Feb 14]. Available from: https://amhonline.amh.net.au/				
	Therapeutic Guidelines. Prevention of neonatal <i>Streptococcus</i> agalactiae (group B streptococcus) disease. In: eTG complete [Internet]. West Melbourne (Victoria): Therapeutic Guidelines; 2019 [cited 2019 Feb 14]. Available from: https://tgldcdp.tg.org.au				
	Therapeutic Guidelines. Syphilis. In: eTG complete [Internet]. West Melbourne (Victoria): Therapeutic Guidelines; 2019 [cited 2019 Feb 14]. Available from: https://tgldcdp.tg.org.au				
	The Royal Women's Hospital. Benzylpenicillin. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2019 [cited 2019 Feb 14]. Available from: https://thewomenspbmg.org.au/				
	Society of Hospital Pharmacists of Australia. Benzylpenicillin. In: Australian Injectable Drugs Handbook [Internet]. [St Leonards, New South Wales]: Health Communication Network; 2019 [cited 2019 Feb 14]. Available from: http://aidh.hcn.com.au				

Keywords:	Benzylpenicillin, benpen, GBS, group B streptococcus, Streptococcus agalactiae, syphilis, syphilus					
Publishing:						
Document owner:	Chief Pharmacist					
Author / Reviewer:	KEMH Pharmacy Department					
Date first issued:	Nov 2014		Version:	3.1		
Last reviewed:	Feb 2019		Next review date:	Feb 2021		
Endorsed by:	Medicines and Therapeutics Committee		Date:	April 2019		
Standards Applicable:	NSQHS Standards: 1 Governance, 3 Infection Control, 4 Medication Safety					

Printed or personally saved electronic copies of this document are considered uncontrolled.

Access the current version from the WNHS website.

For any enquiries relating to this guideline, please email KEMH.PharmacyAdmin@health.wa.gov.au

© Department of Health Western Australia 2019