

ADULT Medication Monograph

MACROGOL with electrolytes

This document should be read in conjunction with this **DISCLAIMER**

Formulary: Unrestricted

Class	Osmotic Laxative				
Presentation	Each powder sachet contains:				
	Macrogol 3350 13.125g				
	Sodium chloride 350.7mg				
	Sodium Bicarbonate 178.5mg				
	Potassium chloride 46.6mg				
Storage	Store at room temperature, below 25°C				
Dose	Constipation				
	Oral:				
	1-3 sachets daily				
	Faecal Impaction				
	Oral:				
	8 sachets daily, taken within 6 hours.				
	Dose may be repeated for up to 3 days.				
Administration	Reconstitute each sachet in a glass containing 125mL of water.				
	For faecal impaction				
	It may be easier to dissolve 8 sachets in 1L of water.				
	If reconstituting larger volumes, store bulk in the fridge (2-8°C) and keep it covered. Discard any solution not used within a 24 hour period.				
Pregnancy	1 st Trimester: Safe to use				
	2 nd Trimester: Safe to use				
	3 rd Trimester: Safe to use				
Breastfeeding	Safe to use				

	T				
Monitoring	Effect is seen between 2-48 hours after dosing				
	Sodium Restriction				
	Each sachet when reconstituted in 125mL of water contains appr 8.1mmol (186mg) of sodium				
	Cardiovascular Disease (e.g. heart failure)				
	Use with caution - may be exacerbated due to fluid and electrolyte disturbances				
Clinical Guidelines	KEMH Clinical Guidelines:				
and Policies	Clinical Practice Guideline – Bowel Care				
	KEMIL Disampassation I & Madiainas Managamant Quidalinas				
	KEMH Pharmaceutical & Medicines Management Guidelines:				
	KEMH Pharmaceutical & Medicines Management Guideline: Medication Administration				
	Australian Medicines Handbook. Macrogol Laxatives. In: Australian				
References	Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2020 [cited 2020 Jun 19]. Available from: https://amhonline.amh.net.au/chapters/gastrointestinal-drugs/laxatives/osmotic-laxatives/macrogol-laxatives				
	MIMS Australia. APO-Macrogol Plus Electrolytes. In: MIMS Online [Internet]. St Leonards (New South Wales): MIMS Australia; 2020 [cited 2020 Jun 19]. Available from: https://www.mimsonline.com.au				
	Therapeutic Guidelines. Functional gastrointestinal disorders: constipation in adults. In: eTG complete [Internet]. West Melbourne (Victoria): Therapeutic Guidelines; 2020 [cited 2020 Jun 19]. Available from: https://tgldcdp.tg.org.au/viewTopic?topicfile=functional-gastrointestinal-disorders&guidelineName=Gastrointestinal#toc_d1e756				
	The Royal Women's Hospital. Macrogols (Polyethylene Glycol (Peg)). In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2020 [cited 2020 Jun 19]. Available from: https://thewomenspbmg-org-au.kelibresources.health.wa.gov.au/medicines/polyethylene-glycol-laxatives/				

Keywords:	Macrogol, Movicol, constipation, bowels, laxative					
Publishing:						
Document owner:	Chief Pharmacist					
Author / Reviewer:	KEMH Pharmacy Department					
Date first issued:	July 2015		Version:	4.0		
Last reviewed:	June 2020		Next review date:	June 2023		
Endorsed by:	Medicines and Therapeutics Committee		Date:	Jan 2018		
Standards Applicable:	NSQHS Standards: 1 Governance, 4 Medication Safety					

Printed or personally saved electronic copies of this document are considered uncontrolled.

Access the current version from the WNHS website.

For any enquiries relating to this guideline, please email KEMH.PharmacyAdmin@health.wa.gov.au

© Department of Health Western Australia 2019