

ADULT Medication Monograph

METHYLENE BLUE (Methylthioninium)

This document should be read in conjunction with this **DISCLAIMER**

Formulary: Unrestricted

SAS Category A (Provedye® requires approval by TGA)

Class	Diagnostic Aid			
Presentation	Prefilled Syringe: 50mg/5mL (Wockhart®)			
Storage	Store at room temperature, below 25°C			
Dose	Diagnostic Aid (Intraoperative)			
	IV Injection: to assist with the visualisation of ureteric jets at cystoscopy			
	Irrigation fluid at hysteroscopy/laparoscopy: to confirm tubal patency			
Compatibility	Wockhart Methylene Blue ® <u>is</u> compatible with sodium chloride 0.9%			
	NB: Different brands of Methylene Blue have a range of compatibilities with Sodium Chloride – Check each product or contact Pharmacy			
Administration	Injection Inject over at least 5 minutes. May be diluted if required			
	Intrauterine irrigation/flush			
	Compatible with Sodium Chloride			
	Flushes Flush with Sodium Chloride			
Monitoring	Monitor for necrosis caused by extravasation			
, 9	Causes staining of skin, saliva, mucous membranes, urine and faeces			
References	Methylene blue (Wockhart®). Product information leaflet. [cited 2021 April 15].			

Keywords:	Methylene blue, Provedye®, Proveblue®, tubal patency, leak test, diagnostic					
Publishing:						
Document owner:	Chief Pharmacist					
Author / Reviewer:	KEMH Pharmacy Department					
Date first issued:	December 2018		Version:	3.0		
Last reviewed:	April 2021		Next review date:	April 2024		
Endorsed by:	Medicines and Therapeutics Committee		Date:	April 2021		
Standards Applicable:	NSQHS Standards: 1 Governance, 4 Medication Safety,					

Printed or personally saved electronic copies of this document are considered uncontrolled.

Access the current version from the WNHS website.

For any enquiries relating to this guideline, please email KEMH.PharmacyAdmin@health.wa.gov.au

© Department of Health Western Australia 2019