



ADULT MEDICATION GUIDELINE

Methylnaltrexone

Scope (Staff): All WNHS Staff

Scope (Area): Obstetrics and Gynaecology

This document should be read in conjunction with the **Disclaimer**.

Quick Links

DoseAdministrationMonitoringPregnancy and Breastfeeding

Formulary: Restricted

Medication Class

Laxative

Presentation

Injection: 12mg/0.6mL vial

Storage

Store at room temperature, below 25°C.

Dose

Opioid induced constipation in palliative care when other regular laxatives inadequate

Subcutaneous injection as single dose:

38 - 62kg: 8mg (0.4mL).

62 - 114kg: 12mg (0.6mL).

Less than 38 or more than 114kg: 0.15mg/kg (or 0.0075mL/kg).

Seek palliative care specialist advice before repeating the injection 24 hours later.

Administration

Refer to the Australian Injectable Drugs Handbook.

Monitoring

Severe or persistent diarrhoea, abdominal pain or cramping.

Pregnancy

1st Trimester: Consider alternative.
2nd Trimester: Consider alternative.
3rd Trimester: Consider alternative.

For more information, please contact **KEMH Obstetric Medicines Information Service**.

Breastfeeding

Consider alternative.

For more information, please contact KEMH Obstetric Medicines Information Service.

Comments

Rapid onset of bowel movement may occur within 30 minutes, with 70% of patients responding within 24 hours. Seek palliative care specialist advice before repeating the injection 24 hours later.

Continuation of (and review of dose) of regular laxatives is mandatory when methylnaltrexone has been used.

Related Policies, Procedures & Guidelines

WNHS Pharmaceutical and Medicines Management Guidelines:

Medication Administration

Palliative Care

References

Australian Medicines Handbook. Methylnaltrexone. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2023 [cited 2023 Feb 07]. Available from: https://amhonline.amh.net.au/

Society of Hospital Pharmacists of Australia. Methylnaltrexone. In: Australian Injectable Drugs Handbook [Internet]. [St Leonards, New South Wales]: Health Communication Network; 2022 [cited 2023 Feb 07]. Available from: http://aidh.hcn.com.au

The Royal Women's Hospital. Methylnaltrexone. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2022 [cited 2023 Feb 12]. Available from: https://thewomenspbmg.org.au/

Keywords	Methylnaltrexone, Constipation, Palliative Care, Relistor					
Document Owner:	Chief Pharmacist					
Author/ Reviewer	KEMH Pharmacy Department					
Version Info:	4.0					
Date First Issued:	07/2015	Last Reviewed:	16/02/2023		Review Date:	16/02/2026
Endorsed by:	Medicines and Therapeutics Committee				Date:	April 2023
NSQHS Standards Applicable:	Std 1: Clinical Governance			Std 5: Comprehensive Care		
	Std 2: Partnering with Consumers			Std 6: Communicating for Safety		
	Std 3: Preventing and Controlling Healthcare Associated Infection			Std 7: Blood Management		
	Std 4: Medication Safety			Std 8: Recognising and Responding to Acute Deterioration		
Printed or personally saved electronic copies of this document are considered uncontrolled.						
Access the current version from WNHS HealthPoint.						

The health impact upon Aboriginal people has been considered, and where relevant incorporated and appropriately addressed in the development of this document (insert ISD Number). (Please refer to the Aboriginal Health Impact Statement and Declaration for Department of Health and Health Service Provider Guidelines – please delete once you have completed this).

This document can be made available in alternative formats on request for a person with a disability.

© Women and Newborn Health Service 2022

Copyright of this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.