

ADULT Medication Monograph

Metoclopramide

This document should be read in conjunction with this **DISCLAIMER**

Formulary: Unrestricted

Class	Dopamine antagonist				
Presentation	Tablets: 10mg				
	Ampoule: 10mg/2mL				
Storage	Tablets: Store below 30°C.				
	Ampoule: Store below 25°C.				
Dose	Nausea and Vomiting				
	Oral, IM or IV injection:				
	>60kg: 10mg three times daily as needed.				
	Maximum 30mg or 0.5mg/kg in 24 hours (whichever is less).				
	NB: Should not be used for more than 5 days at a time.				
	Gastric Stasis				
	Oral, IM or IV injection:				
	>20 years: 5-10mg three times daily. Maximum 30mg in 24 hours.				
	NB: Should not be used for more than 5 days at a time.				
Administration	Oral Take with or without food				
	<u>Intramuscular</u>				
	Inject into a large muscle				
	IV Injection				
	IV Injection Cive over at least 2 minutes to minimise transient egitation and				
	Give over at least 3 minutes to minimise transient agitation and restlessness.				
	IV Infusion				
	Step 1 Dilution: Dilute the dose in at least 50 mL of compatible fluid.				
	Step 2 Administration: Infuse over 15 minutes.				

Pregnancy	1 st Trimester: Safe to use				
	2 nd Trimester: Safe to use				
	3 rd Trimester: Safe to use				
Breastfeeding	Considered safe to use				
Clinical Guidelines and Policies	KEMH Clinical Guidelines:				
	Obstetrics: Caesarean birth				
	Obstetrics: Pregnancy care: First trimester complications				
	Obstetrics: Ovarian hyperstimulation syndrome				
	Obstetrics: Antenatal Care: Minor symptoms and disorders in pregnancy				
	Gynaecology: Palliative Care				
	Neonates: <u>Gastro-Oesophageal Reflux</u>				
	KEMH Pharmaceutical and Medicines Management Guidelines:				
	KEMH Pharmaceutical & Medicines Management Guideline: <u>Medication Administration</u>				
References	Australian Medicines Handbook. Metoclopramide. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2020 [cited 2020 Mar 25]. Available from: https://amhonline.amh.net.au/				
	MIMS Australia. Pramin. In: MIMS Online [Internet]. St Leonards (New South Wales): MIMS Australia; 2020 [cited 2020 Mar 25]. Available from: https://www.mimsonline.com.au				
	The Royal Women's Hospital. Metoclopramide. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2020 [cited 2020 Mar 25]. Available from: https://thewomenspbmg.org.au/				
	Society of Hospital Pharmacists of Australia. Metoclopramide. In: Australian Injectable Drugs Handbook [Internet]. [St Leonards, New South Wales]: Health Communication Network; 2020 [cited 2020 Mar 25]. Available from: http://aidh.hcn.com.au				

Keywords:	Hyperemesis, nausea, vomiting, gastric stasis, metoclopramide				
Publishing:					
Document owner:	Chief Pharmacist				
Author / Reviewer:	KEMH Pharmacy Department				
Date first issued:	Dec 2014		Version:	4.0	
Last reviewed:	April 2020		Next review date:	Apr 2023	
Endorsed by:	Medicines and Therapeutics Committee		Date:		
Standards Applicable:	NSQHS Standards:				
	1 Governance, 4 Medication Safety				

Printed or personally saved electronic copies of this document are considered uncontrolled.

Access the current version from the WNHS website.

For any enquiries relating to this guideline, please email KEMH.PharmacyAdmin@health.wa.gov.au

© Department of Health Western Australia 2020