

Government of Western Australia North Metropolitan Health Service Women and Newborn Health Service



ADULT MEDICATION GUIDELINE PROBENECID Scope (Staff): All WNHS Staff Scope (Area): Obstetrics and Gynaecology This document should be read in conjunction with the Disclaimer.

| Quick Links | | | | | | | |
|-----------------------------|----------------|------------|--------------------------------|--|--|--|--|
| Dose | Administration | Monitoring | Pregnancy and Breastfeeding | | | | |
| Restrictions | | | | | | | |
| Formulary: Unrestricted | | | | | | | |
| Medication Class | | | | | | | |
| Antibiotic thereasy adjunct | | | | | | | |

Antibiotic therapy adjunct

Antigout agent

Antihyperuricaemic

Presentation

Tablet: 500mg

Storage

Store at room temperature, below 30°C

Dose

Antibiotic therapy, adjunct

• Probenecid is indicated for the elevation and prolongation of plasma levels of beta-lactam antibiotics, irrespective of the route of administration of the antibiotic.

Gonococcal infection

If susceptibility to penicillin is confirmed, or in areas where penicillin-resistant N. gonorrhoeae is less common (mainly in some remote areas of northern and central Australia), to treat asymptomatic genital gonococcal infection, use:

Oral:

1g when used in combination with amoxicillin 3g and azithromycin 1g (all as single doses)

Administration

<u>Oral</u>

Take with food to reduce stomach upset. Ensure adequate fluid intake (to prevent kidney stones)

Monitoring

Risk of uric acid kidney stones

Check renal function and complete blood count regularly

Pregnancy

1st Trimester: Monitoring required

2nd Trimester: Monitoring required

3rd Trimester: Monitoring required

Breastfeeding

Considered safe to use

Comments

Prohibited in elite sports as it may be used to mask detection of banned substances

Related Policies, Procedures & Guidelines

Department of Health WA Policies:

Silver Book- STI/BBV Management Guidelines WA

WNHS Policies:

Antimicrobial Stewardship (AMS) policy

WNHS Clinical Practice Guidelines:

Sexually Transmitted Infections (STI)

References

Australian Medicines Handbook. Probenecid. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2021 [cited 2021 Sep 21]. Available from: https://amhonline.amh.net.au/ Therapeutic Guidelines. Approach to *Neisseria gonorrhoeae*. In: eTG complete [Internet]. West Melbourne (Victoria): Therapeutic Guidelines; 2021 [cited 2021 Sep 21]. Available from: https://tgldcdp-tg-org-au.kelibresources.health.wa.gov.au/etgAccess

The Royal Women's Hospital. Probenecid. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2021 [cited 2021 Sep 21]. Available from: https://thewomenspbmg.org.au/

Sexual Health and Blood-borne Virus Program. Notifiable infections: gonorrhoea. Silver book guidelines for managing sexually transmitted infections and blood-borne viruses [Internet]. Perth (WA): Government of Western Australia Department of Health; 2015 [updated 2020] cited 2021 Sep 21]. Available from: http://ww2.health.wa.gov.au/Silver-book/Notifiableinfections/Gonorrhoea.

| Keywords | Probenecid, gonococcal infection, gonorrhoea | | | | | | | | |
|---|---|----------------|----------|--|--------------|------------|--|--|--|
| Document Owner: | Chief Pharmacist | | | | | | | | |
| Author/ Reviewer | KEMH Pharmacy Department | | | | | | | | |
| Version Info: | 4.0 | | | | | | | | |
| Date First Issued: | July 2015 | Last Reviewed: | Sep 2021 | | Review Date: | Nov 2024 | | | |
| Endorsed by: | Medicines and Therapeutics Committee | | | | Date: | 02/11/2021 | | | |
| NSQHS Standards | Std 1: Clinical Governance | | | Std 5: Comprehensive Care | | | | | |
| Applicable: | Std 2: Partnering with Consumers | | | Std 6: Communicating for Safety | | | | | |
| | Std 3: Preventing and Controlling Healthcare Associated Infection | | | Std 7: Blood Management | | | | | |
| | Std 4: Medication Safety | | | Std 8: Recognising and Responding to Acute Deterioration | | | | | |
| Printed or personally saved electronic copies of this document are considered uncontrolled. | | | | | | | | | |
| Access the current version from WNHS HealthPoint. | | | | | | | | | |

The health impact upon Aboriginal people has been considered, and where relevant incorporated and appropriately addressed in the development of this document (insert ISD Number). (Please refer to the Aboriginal Health Impact Statement and Declaration for Department of Health and Health Service Provider Guidelines – please delete once you have completed this).

This document can be made available in alternative formats on request for a person with a disability. © Women and Newborn Health Service 2021 Copyright of this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.