



ADULT MEDICATION GUIDELINE TOBRAMYCIN Scope (Staff): All WNHS Staff Obstetrics and Gynaecology

This document should be read in conjunction with the Disclaimer.

Quick Links

Scope (Area):

Pregnancy and **Administration Monitoring** <u>Dose</u> Breastfeeding

Restrictions

Formulary: Highly Restricted

HIGH RISK Medication A

Medication Class

Drug Class

Presentation

Vial: 80mg/2mL

Also contain sodium metabisulfite, disodium edetate, sulfuric acid and/or sodium hydroxide. Pfizer brand also contains phenol.

Storage

DBL: store below 25 C. Protect from light. Pfizer: store below 30 C. Protect from light.

Dose

As per Gentamicin Dosing and Monitoring

Administration

Refer to the Australian Injectable Drugs Handbook

Monitoring

Refer to Gentamicin Dosing and Monitoring

Pregnancy

1st Trimester: Monitoring required
2nd Trimester: Monitoring required
3rd Trimester: Monitoring required

Breastfeeding

Considered safe to use

Comments

Tobramycin is inactivated by penicillin and cephalosporin antibiotics.

- Do not mix in the same injection or infusion solution.
- Administer at separate sites if possible.
- In patients with normal renal function where it is not practical or possible to administer separately, flush the line well before and after giving each drug.

Tobramycin for injection may contain excipients that are not appropriate for inhalation. Use the products that are formulated for inhalation in nebulisers.

See Practice points from Aminoglycosides in AMH

Related Policies, Procedures & Guidelines

WNHS Clinical Practice Guidelines:

Gentamicin Dosing and Monitoring

WNHS Pharmaceutical and Medicines Management Guidelines:

HIGH RISK Medication

References

Australian Medicines Handbook. Benzathine penicillin. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2017 [cited 2017 Apr 12]. Available from: https://amhonline.amh.net.au/

Therapeutic Guidelines. Acute pain: perioperative. In: eTG complete [Internet]. West Melbourne (Victoria): Therapeutic Guidelines; 2017 [cited 2017 Apr 12]. Available from: https://tgldcdp-tg-org-au.kelibresources.health.wa.gov.au/etgAccess

The Royal Women's Hospital. Folic acid. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2016 [cited 2017 Apr 12]. Available from: https://thewomenspbmg.org.au/

Tobramycin					
Chief Pharmacist					
KEMH Pharmacy Department					
4.0					
2017	Last Reviewed:	Feb 2022		Review Date:	Feb 2025
Medicines and Therapeutics Committee				Date:	01/03/2022
Std 1: Clinical Governance			Std 5: Comprehensive Care		
Std 2: Partnering with Consumers			Std 6: Communicating for Safety		
Std 3: Preventing and Controlling Healthcare Associated Infection			Std 7: Blood Management		
Std 4: Medication Safety			Std 8: Recognising and Responding to Acute Deterioration		
Printed or personally saved electronic copies of this document are considered uncontrolled.					
	Chief Pharmacis KEMH Pharmacy 4.0 2017 Medicines and T Std 1: Cl Std 2: P Std 3: P Healthcare Asso Std 4: Mo	Chief Pharmacist KEMH Pharmacy Department 4.0 2017 Last Reviewed: Medicines and Therapeutics Comm Std 1: Clinical Governance Std 2: Partnering with Const Std 3: Preventing and Cont Healthcare Associated Infection Std 4: Medication Safety Dersonally saved electronic copies	Chief Pharmacist KEMH Pharmacy Department 4.0 2017 Last Reviewed: Feb 2022 Medicines and Therapeutics Committee Std 1: Clinical Governance Std 2: Partnering with Consumers Std 3: Preventing and Controlling Healthcare Associated Infection Std 4: Medication Safety Dersonally saved electronic copies of this description.	Chief Pharmacist KEMH Pharmacy Department 4.0 2017	Chief Pharmacist KEMH Pharmacy Department 4.0 2017

The health impact upon Aboriginal people has been considered, and where relevant incorporated and appropriately addressed in the development of this document (insert ISD Number). (Please refer to the Aboriginal Health Impact Statement and Declaration for Department of Health and Health Service Provider Guidelines – please delete once you have completed this).

This document can be made available in alternative formats on request for a person with a disability.

© Women and Newborn Health Service 2022

Copyright of this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.