



11/10/2021

Please be advised, this document is under comprehensive review.
Current content and practices outlined within this document must be followed until the review is complete.

CLINICAL GUIDELINE

Hepatitis C Virus (HCV): Care of the Infant Born to HCV Positive Women

Scope (Staff):	Nursing and Medical Staff
Scope (Area):	KEMH Postnatal Wards

This document should be read in conjunction with this [DISCLAIMER](#)

This guideline should be used in conjunction with its respective Obstetrics and Midwifery Clinical Guidelines - [Antenatal Care: Hepatitis B in Pregnancy](#).
Refer to Neonatal Clinical Guideline - [Hepatitis C Virus \(HCV\)](#) for additional information.

Key Points

- Factors that increase risk for perinatal transmission of HCV include:
 - Maternal detection of HCV RNA by the PCR test^{1, 2}
 - Peripheral blood mononuclear cell infection¹ e.g. HIV co-infection²
 - Prolonged rupture of membranes for more than 6 hours¹
 - Obstetric procedures or intrapartum exposure to maternal blood infected by HCV e.g. vaginal or perineal lacerations¹
 - Females are twice as likely to be infected as males².
- Approximately 5% of chronic HCV positive pregnant women will transmit the virus to the infant.³ In most cases vertical transmission is confined to women with detectable HCV RNA.²
- Maternal co-infection with HCV and HIV (in women untreated with HIV antiviral agents) is associated with increased risk of perinatal transmission.¹
- Breastfeeding is recommended provided the nipples are not damaged or bleeding. Milk should be expressed and discard until nipples are healed.⁴
- The neonate should be bathed to remove maternal body secretions prior to administering intramuscular injections.⁴
- HCV infection acquired from perinatal transmission progresses to chronic HCV in 80% of cases, with very few children clearing it spontaneously.¹
- Diagnosis of perinatal transmission is made on anti-HCV antibody testing, and/or by detecting HCV RNA on peripheral blood by performing a PCR test.^{1, 5}

Hepatitis C Virus (HCV): Care of the Infant Born to HCV Positive Women

- Cord blood testing of HCV is not recommended as it can yield false-positive or false-negative results.⁵
- It is recommended that a child at risk of perinatal transmission exposure to HCV should be tested for HCV antibodies at 18 months of age. Testing prior to this time is limited due to passive transfer of maternal antibodies.^{1, 3, 5}
- Should diagnosis of vertical transmission of HCV be necessary prior to 18 months of age (e.g. parental request) then testing of infant serum HCV RNA can be obtained when the infant is between 2 and 6 months of age.⁶ A sample is collected on two occasions at least 3 - 4 months apart to confirm diagnosis.^{2, 5}
- Hepatitis B virus (HBV) vaccination is recommended within 12 hours of birth as these neonates are at higher risk of infection, and if a child is infected with HCV they may be more prone to severe infection if they contract HBV.³
- The paediatrician shall generate a [letter to the GP](#) with the recommended follow-up regimen for neonates of HCV infected mothers

Related CAHS internal policies, procedures and guidelines

Obstetrics and Midwifery Clinical Guidelines

- [Antenatal Care: Hepatitis B in Pregnancy](#)

Neonatal Clinical Guideline

- [Hepatitis C Virus \(HCV\)](#)

References and related external legislation, policies, and guidelines



1. Indolfi G, Resti M. Perinatal Transmission of Hepatitis C Virus Infection. **Journal of Medical Virology**. 2009;81:836-4.
2. Davison SM, Mieli-Vergani G, Sira J, et al. Perinatal hepatitis C virus infection: diagnosis and management. **Archives of Disease in Childhood**. 2006;91(781-785).
3. Canadian Paediatric Society. Position Statement. Vertical transmission of the hepatitis C virus: Current knowledge and issues. **Paediatric Child Health**. 2008;13(6):529-34.
4. The Royal Australian and New Zealand College of Obstetricians and Gynaecologists. C-Gen 4 Hepatitis C. **RANZCOG College statement**. 2009.
5. Valladares G, Sjogren MH, Chacaltana A. The management of HCV-infected pregnant women. **Annals of Hepatology**. 2010;9(Suppl.):S92-7.
6. Jain S, Goharkhay N, Saade G, et al. Hepatitis C in Pregnancy. **American Journal of Perinatology**. 2007;24:251-6.

Useful resources (including related forms)

[GP Referral Letter](#)

Hepatitis C Virus (HCV): Care of the Infant Born to HCV Positive Women

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Neonatology		
Reviewer / Team:	Neonatal Coordinating Group		
Date First Issued:	August 2011	Last Reviewed:	25 th July 2017
Amendment Dates:		Next Review Date:	25th July 2020
Approved by:	Neonatal Coordinating Group	Date:	26 th September 2017
Endorsed by:	Neonatal Coordinating Group	Date:	
Standards Applicable:	NSQHS Standards:  		
Printed or personally saved electronic copies of this document are considered uncontrolled			